

## Chemokines, Novel Lymphocyte Subsets and Lymphocyte Trafficking

Stanford University Medical Center, Stanford, California  
and Veterans Administration, Palo Alto, California

Chemokines control lymphocyte trafficking and microenvironmental positioning. I will discuss recent observations that 1) reveal the role of constitutive tissue-specific epithelial-cell chemokines in effector and antibody secreting cell homing to mucosal and cutaneous sites; 2) use chemokine and homing receptor expression patterns to identify novel T cell subsets, including separable CD4 subsets responsible for B help and DTH; 3) demonstrate that specific homing properties are induced rapidly by local microenvironments during the naive to effector/memory transition in lymphoid tissues; and 4) redefine the rules and significance of chemokine receptor association with Th1 and Th2 cells. These novel findings will be considered in the context of the combinatorial control of lymphocyte trafficking, and its intimate interrelationship with lymphocyte differentiation and function.

Campbell, D. J., Kim, C. H., and Butcher, E. C. (2001). "Separable effector T cell populations specialized for B cell help or tissue inflammation." *Nat Immunol*, 2(9), 876-81.

Kim, C. H., Rott, L., Kunkel, E. J., Genovese, M. C., Andrew, D. P., Wu, L., and Butcher, E. C. (2001a). "Rules of chemokine receptor association with T cell polarization in vivo." *J Clin Invest*, 108(9), 1331-9.

Kim, C. H., Rott, L. S., Clark-Lewis, I., Campbell, D. J., Wu, L., and Butcher, E. C. (2001b). "Subspecialization of CXCR5+ T cells: B helper activity is focused in a germinal center-localized subset of CXCR5+ T cells." *J Exp Med*, 193(12), 1373-81.

Pan, J., Kunkel, E. J., Gossler, U., Lazarus, N., Langdon, P., Broadwell, K., Vierra, M. A., Genovese, M. C., Butcher, E. C., and Soler, D. (2000). "A novel chemokine ligand for CCR10 and CCR3 expressed by epithelial cells in mucosal tissues." *J Immunol*, 165(6), 2943-9.

Reiss, Y., Proudfoot, A. E., Power, C. A., Campbell, J. J., and Butcher, E. C. (2001). "CC chemokine receptor (CCR)4 and the CCR10 ligand cutaneous T cell-attracting chemokine (CTACK) in lymphocyte trafficking to inflamed skin." *J Exp Med*, 194(10), 1541-7.