

Adoptive T cell receptor (TCR) gene transfer of HPV16 HLA-A*0201 Specific TCR to Human T Cells

Eiben GL¹, Nishimura MI² and Kast WM³

(1) Cardinal Bernardin Cancer Center, Loyola University Chicago, Maywood IL 60153.
(2) Dept. of Surgery, University of Chicago, Chicago, IL 60637. (3) Norris Comprehensive Cancer Center, University of Southern California, Los Angeles, CA 90089.

A T cell mediated response against the early proteins of HPV may offer considerable therapeutic potential against cervical cancer. Recently, TCR gene transfer has been proposed as a strategy to provide anti-tumor immunity to cancer patients. It has been shown that transferring the TCR genes from tumor reactive T cell clones to peripheral blood lymphocytes can redirect their specificity to recognize the appropriate tumor antigens. Therefore, we have identified and cloned the α and β chains from the TCR of two CTL clones reactive with the HLA-A*0201 restricted epitopes from HPV16 E6 29-38 and E7 11-20. By placing the α and β chains of these TCR into retroviral vectors, we can transfer the specific TCR to human T cell lines and PBL and detect the receptor by staining with TCR specific V β subfamily antibodies and HPV16 specific peptide/MHC tetramers. Furthermore, the recipient T cells can elicit peptide specific immune responses as detected by cytokine release and lysis assays. The ability to adoptively transfer genetically engineered tumor reactive T cells to a patient may represent an effective anti-tumor therapy against cervical cancer and this study may help pave the way for clinical evaluation of this treatment strategy.